

Neutrino Oscillation in Matter and Parameters s_{13}, δ_{CP}

Leonard S. Kisslinger^a, Ernest M. Henley^b, and Mikkell B. Johnson^c

^aDepartment of Physics, Carnegie Mellon University, Pittsburgh, PA 15213

^bDepartment of Physics, University of Washington, Seattle, WA 98195

^cLos Alamos National Laboratory, Los Alamos, NM 87545

March 1, 2013

PACS Indices:11.30.Er,14.60.Lm,13.15.+g

Abstract

We estimate the dependence of ν_μ to ν_e conversion on parameters θ_{13} and δ_{CP} for several experimental facilities studying neutrino oscillations. We use the S-Matrix theory to estimate $\bar{\nu}_e$ disappearance and compare estimates based on an older theory being used to extract θ_{13} from the Double Chooz, Daya Bay, and RENO data, to assist in extracting an accurate value for θ_{13} from these projects. We use values of θ_{13} within known limits, and estimate the dependence of ν_μ - ν_e CP violation (CPV) probability on δ_{CP} in order to suggest new experiments to measure CPV for neutrinos moving in matter.

1 Introduction

In our present work we study ν_μ to ν_e neutrino conversion, $\bar{\nu}_e$ disappearance, and CP violation (CPV) measurements using the S-Matrix method for neutrino oscillations. The study of CP violation is essential for understanding weak interactions. Almost half a century ago CP violation in weak interactions was found in the decay of K_L^0 into $\pi^+ + \pi^-$ [1] and $2\pi^0$ [2], with branching ratios of the order of .001. The decay $K_L^0 \rightarrow \pi^0 + \nu + \bar{\nu}$ is almost entirely CP violating [3] but requires accurate determination of the CKM matrix [4] and

accurate measurements. See Ref[5] for a review of this experiment and references. There have many other studies of CP asymmetries in weak decays: see Ref[6] for a recent study of $\bar{B}$ radiative decay with references to earlier work on CP violation in various weak decays.

In recent years there have been a number of experimental studies of neutrino oscillations using neutrino beams from accelerators and reactors, and important objectives of these experiments are the measurements of ν_μ to ν_e conversion and CPV. The first study in our present work is an estimate the ν_μ to ν_e conversion probability using parameters for the baseline and energy corresponding to MiniBooNE[7], JHF-Kamioka [8], MINOS[9], and CHOOZ[10]-Double Chooz[11], which are on-going projects, although the CHOOZ project does not have a beam of ν_μ neutrinos.

There have been many recent studies of CP and T symmetries via neutrino oscillations for future facilities, e.g., see Refs[12, 13], which also give references to earlier publications, and the ISS report[14] on future neutrino facilities. The two main parameters of interest in the present work are δ_{CP} , which is essentially unknown, and the angle θ_{13} . One possible future facility for studying CPV and the δ_{CP} parameter is the LBNE Project, where neutrino beams produced at Fermilab would have a baseline of $L \simeq 1200$ km, being detected with deep underground detectors[15, 16]. With the methods used in the present work, described below, predictions of CPV with the baseline and energies of the LBNE Project have recently been made for δ_{CP} from 90 to 0 degrees[17].

Although the angle θ_{13} is not well known, there are limits on its value. Recently the T2K collaboration [18] published limits, and found a best-fit value of $\theta_{13} \simeq 11$ degrees, or $\sin\theta_{13} \equiv s_{13} \simeq 0.19$, which is one of the values we use in the present work on neutrino transition probability in Sec. 2.

The angle θ_{13} will be measured by the Daya Bay experiment[19, 20] in China, the Double Chooz project[21, 22] in France, and RENO[23] in Korea, via $\bar{\nu}_e$ disappearance. A very recent result from the Daya Bay project[24] concludes that $s_{13} \simeq 0.15$, and in our study (Sec. 3) of $\bar{\nu}_e$ disappearance we use the the Daya Bay parameters to test the theory. Our results will also be useful for the Double Chooz and RENO projects.

Finally, in Sec 4, using the expected range of values for θ_{13} , we estimate CPV for $\mu - e$ neutrino oscillation for the entire range of δ_{CP} to help in the planning for future CPV experiments.

A major complication for the determination of T, CP, and CPT violation is the interaction of neutrinos with matter as they travel along the baseline. These matter effects have been studied by a number of theorists. See, e.g., Refs[25, 26, 27]. One main objective of the present research is to estimate matter effects for the MiniBooNE, JHF-Kamioka, MINOS, and CHOOZ facilities, to help find the values of θ_{13} and δ_{CP} .

For the basic interactions, which are CPT invariant for local theories, CP and T violation have the same magnitude. With matter effects T and CP are not directly related. Our present research is an extension of our recent work on T reversal violation[28]. In that study we used the formalism of Ref[29] for $\nu_e \leftrightarrow \nu_\mu$ TRV, and that of Ref[30] for $\nu_e \rightarrow \nu_\mu$ conversion probability to calculate the effects of neutrinos moving through matter. In the present work we use the notation and formalism of Jacobson and Ohlsson[31], who studied possible matter effects for CPT violation.

CP violation in the $a - b$ sector is given by the transition probability, denoted by $\mathcal{P}(\nu_a \rightarrow \nu_b)$, for a neutrino of flavor a to convert to a neutrino of flavor b ; and similarly for antineutrinos $\bar{\nu}_a, \bar{\nu}_b$. The CPV probability differences are defined as

$$\Delta\mathcal{P}_{ab}^{CP} = \mathcal{P}(\nu_a \rightarrow \nu_b) - \mathcal{P}(\bar{\nu}_a \rightarrow \bar{\nu}_b) . \quad (1)$$

In our present work we study $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ and $\mathcal{P}(\nu_\mu \rightarrow \nu_e) - \mathcal{P}(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$ since the neutrino beams at MiniBooNE, JHF-Kamioka, and MINOS, are muon or anti-muon neutrinos. We then calculate $\bar{\nu}_e$ disappearance probability for Daya Bay baseline and energy, comparing our S-Matrix theory to the formula used by Daya Bay, Double Chooz and RENO[19, 20, 21, 22, 23] .

2 Transition Probability $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ and $\bar{\nu}_e$ Disappearance

In this section we review ν_μ to ν_e oscillation probability derived from standard S-matrix theory and then compare the probability of $\bar{\nu}_e$ disappearance derived from this theory to that used by the Double Chooz experimental project[21, 22].

2.1 $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ Derived Using S-Matrix Theory

In this subsection we review the derivation of the probability of a muon neutrino to convert to an electron neutrino, $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$, using the notation of Ref[31]. We then make an estimate of the transition probabilities for sample accelerator and reactor experiments. Although at the present time no experiments for CPV are possible, this can serve as a basis for future experiments. In the next section we give somewhat more accurate calculations for CPV for the same set of experimental facilities.

As in Refs[29, 31] we use the time evolution matrix, $S(t, t_0)$ to derive the transition probabilities. For neutrino oscillations the initial neutrino beam is emitted at time t_0 , usually taken as $t_0 = 0$, and the neutrino or converted neutrino is detected at baseline length $= L$ at time $= t$. Since the

neutrinos move with a velocity near that of the speed of light, at the end of our derivation we take $t - t_0 \rightarrow L$, with the units $c=1$.

Given the Hamiltonian, $H(t)$, for neutrinos, the neutrino state at time t is obtained from the state at time t_0 from the matrix, $S(t, t_0)$, by

$$|\nu(t)\rangle = S(t, t_0)|\nu(t_0)\rangle \quad (2)$$

$$i\frac{d}{dt}S(t, t_0) = H(t)S(t, t_0) . \quad (3)$$

Neutrinos (and antineutrinos) are produced as ν_a , where a is the flavor, $a = e, \mu, \tau$. However, neutrinos of definite masses are ν_α , with $\alpha = 1, 2, 3$. The two forms are connected by a 3 by 3 unitary transformation matrix, U : $\nu_a = U\nu_\alpha$, where ν_a, ν_α are 3x1 column vectors and U is given by ($\sin\theta_{ij} \equiv s_{ij}$)

$$U = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{CP}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{CP}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{CP}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{CP}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{CP}} & c_{23}c_{13} \end{pmatrix} ,$$

similar to the CKM matrix for quarks. We use the best fit value[12] $s_{23} = 0.707$. θ_{13} and $s_{12} = 0.56$, $c_{12} = 0.83$. We use $s_{13} = 0.19$ and $s_{13}=0.095$, as discussed above, to determine the dependence of $\nu_\mu \rightarrow \nu_e$ conversion, and CPV on this parameter. We calculate the dependence of $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ and $\Delta\mathcal{P}_{\mu e}^{CP}$ on δ_{CP} , as discussed below. In the vacuum the $S(t, t_0)$ is obtained from

$$S_{ab}(t, t_0) = \sum_{j=1}^3 U_{aj} \exp(iE_j(t-t_0)) U_{bj}^* . \quad (4)$$

Since neutrino beams in neutrino oscillation experiments travel through matter, and the main neutrino-matter is scattering from electrons, we must include potential, $V = \sqrt{2}G_F n_e$, for neutrino electron scattering in the earth: where G_F is the universal weak interaction Fermi constant, and n_e is the density of electrons in matter. Using the matter density $\rho=3$ gm/cc, the neutrino-matter potential is $V = 1.13 \times 10^{-13}$ eV.

The transition probability $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ is obtained from S_{12} :

$$\mathcal{P}(\nu_\mu \rightarrow \nu_e) = |S_{12}|^2 = \text{Re}[S_{12}]^2 + \text{Im}[S_{12}]^2 , \quad (5)$$

with

$$S_{12} = c_{23}\beta - is_{23}ae^{-i\delta_{CP}}A \quad (6)$$

$$a = s_{13}(\Delta - s_{12}^2\delta) \quad (7)$$

$$\delta = \delta m_{12}^2/(2E) \quad (8)$$

$$\Delta = \delta m_{13}^2/(2E) \quad (9)$$

$$A \simeq f(t)I_\alpha * \quad (10)$$

$$I_\alpha * = \int_0^t dt' \alpha^*(t') f(t') \quad (11)$$

$$\alpha(t) = \cos\omega t - i\cos 2\theta \sin\omega t \quad (12)$$

$$f(t) = e^{-i\bar{\Delta}t} \quad (13)$$

$$2\omega = \sqrt{\delta^2 + V^2 - 2\delta V \cos(2\theta_{12})} \quad (14)$$

$$\beta = -i\sin 2\theta \sin\omega L \quad (15)$$

$$\bar{\Delta} = \Delta - (V + \delta)/2 \quad (16)$$

$$\sin 2\theta = s_{12}c_{12}\frac{\delta}{\omega}, \quad (17)$$

where the neutrino mass differences are $\delta m_{12}^2 = 7.6 \times 10^{-5}(eV)^2$ and $\delta m_{13}^2 = 2.4 \times 10^{-3}(eV)^2$. Note that $\delta \ll \Delta$, and $t \rightarrow L$ for $v_\nu \simeq c$. From Eqs.(5 to 16):

$$\begin{aligned} Re[S_{12}] &= s_{23}a[\cos(\bar{\Delta}L + \delta_{CP})Im[I_{\alpha*}] - \sin(\bar{\Delta}L + \delta_{CP})Re[I_{\alpha*}]] \\ Im[S_{12}] &= -c_{23}\sin 2\theta \sin\omega L - s_{23}a[\cos(\bar{\Delta}L + \delta_{CP})Re[I_{\alpha*}] \\ &\quad + \sin(\bar{\Delta}L + \delta_{CP})Im[I_{\alpha*}]] \end{aligned} \quad (18)$$

Using $\delta, \omega \ll \Delta$ one can show that

$$\begin{aligned} Re[I_{\alpha*}] &\simeq \frac{\sin\bar{\Delta}L}{\bar{\Delta}} \\ Im[I_{\alpha*}] &\simeq \frac{1 - \cos\bar{\Delta}L}{\bar{\Delta}}. \end{aligned} \quad (19)$$

From Eqs(18,19) we find

$$\begin{aligned} \mathcal{P}(\nu_\mu \rightarrow \nu_e) &\simeq (c_{23}s_{12}c_{12}(\delta/\omega)\sin\omega L)^2 + 2(s_{23}s_{13})^2(1 - \cos\bar{\Delta}L) \\ &\quad + 2s_{13}s_{12}c_{12}s_{23}c_{23}(\delta/\omega)\sin\omega L \\ &\quad (\cos(\bar{\Delta}L + \delta_{CP})\sin\bar{\Delta}L + \sin(\bar{\Delta}L + \delta_{CP})(1 - \cos\bar{\Delta}L)). \end{aligned} \quad (20)$$

We use $s_{13}=0.19$ and 0.095 to show the effect of s_{13} .




Figure 1: The ordinate is $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ for MINOS($L=735$ km), MiniBooNE($L=500$ m), JHF-Kamioka($L=295$ km), and CHOOZ($L=1.03$ km). Solid curve for $s_{13}=0.19$ and dashed curve for $s_{13}=0.095$. The curves are almost independent of δ_{CP} .

From Eq(20) we obtain the results for $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ shown in Fig.1. These results can provide guidance for future experiments on CPV via $\nu_\mu \leftrightarrow \nu_e$ oscillation. Note that in Ref[8] $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ was calculated for the 295 km JHF-Kamioka project for $E=0-2$ GeV, and our calculation based on the theory developed in Refs.[31, 29], finds $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ is in agreement for $E=.4-1.0$ GeV with this earlier estimate. We calculated $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ for δ_{CP} from $-\pi/2$ to $\pi/2$, and the results are almost independent of δ_{CP} . The results for CHOOZ are shown in preparing for the following subsection on $\bar{\nu}_e$ disappearance, even though Double Chooz, Daya Bay, and RENO projects have beams of $\bar{\nu}_e$ rather than ν_μ neutrinos.

3 $\bar{\nu}_e$ Disappearance Derived Using S-Matrix Theory Compared to Daya Bay Evaluation

In this subsection we derive $\bar{\nu}_e$ disappearance, $\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e)$, defined as

$$\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e) = 1 - \mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_\mu) - \mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_\tau), \quad (21)$$

using the S-matrix method (see previous subsection), and compare it to the expression for $\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e)$ used by the Daya Bay, Double Chooz, and RENO, which is (see, e.g., Refs[11, 21])

$$\mathcal{P}^{\mathcal{DB}}(\bar{\nu}_e \rightarrow \bar{\nu}_e) \simeq 1 - 4(s_{13}c_{13})^2 \sin^2\left(\frac{\Delta L}{2}\right) \quad (22)$$

where $\Delta \equiv \delta m_{13}^2/(2E)$, Eq(8), and $s_{13}, c_{13} = \sin\theta_{13}, \cos\theta_{13}$. A third term with a factor of $\sin^2(\delta L/2)$ [22] was dropped[11, 21] because $\delta m_{12}^2 \ll \delta m_{13}^2$ and $\sin^2(\delta L/2) \ll \sin^2(\Delta L/2)$ for Daya Bay baseline $L=1.9$ km and energy $E=4$ MeV.

In the S-matrix method (see Ref[31]) the probability of $\bar{\nu}_e$ oscillation to $\bar{\nu}_\mu$ and $\bar{\nu}_\tau$ is given by (see, e.g., Ref[31])

$$\begin{aligned} \mathcal{P}^{\mathcal{SM}}(\bar{\nu}_e \rightarrow \bar{\nu}_\mu) &= |\bar{S}_{21}|^2 \\ \mathcal{P}^{\mathcal{SM}}(\bar{\nu}_e \rightarrow \bar{\nu}_\tau) &= |\bar{S}_{31}|^2. \end{aligned} \quad (23)$$

We take $\delta_{CP} = 0$, since $|S_{12}|^2$ is essentially independent of δ_{CP} , so $A = C$ (see Ref[29] for definition and proof). Therefore $|\bar{S}_{21}|^2 = |S_{12}(V \rightarrow -V)|^2$, and $|\bar{S}_{31}|^2 = |S_{12}|^2(V \rightarrow -V, c_{23} \rightarrow s_{23}, s_{23} \rightarrow -c_{23})$. Using $s_{23}^2 = c_{23}^2 = 1/2$ we find

$$\mathcal{P}^{\mathcal{SM}}(\bar{\nu}_e \rightarrow \bar{\nu}_e) = 1 - [(.46\delta \sin\bar{\omega}L/\bar{\omega})^2 + 2(s_{13})^2(1 - \cos\bar{\Delta}L)], \quad (24)$$

with $\bar{\Delta} = \Delta + (V - \delta)/2$ and $2\bar{\omega} = \sqrt{\delta^2 + V^2 + 2\delta V \cos(2\theta_{12})}$

Fig. 2 for $L=1.9$ km and Fig. 3 for $L=10$ km are discussed below.




Figure 2: $\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e)$. For $s_{13} = .15$ (a) $\mathcal{P}^{DB}$ and (b) $\mathcal{P}^{SM}$

From Fig. 2 the ratios $R1, R2$, of $1 - \mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e)$ for $\mathcal{P}^{DB}$ (Eq(22)) to $\mathcal{P}^{SM}$ (Eq(24)) for $s_{13} = .15$, and for $s_{13} = .15$ for $\mathcal{P}^{DB}$ and $s_{13} = .147$ for $\mathcal{P}^{SM}$, for $E \simeq 4.0\text{MeV}$ and $L=1.9\text{ km}$ are

$$\begin{aligned} R1 &= \frac{1 - \mathcal{P}^{DB}(s_{13} = .15)}{1 - \mathcal{P}^{SM}(s_{13} = .15)} = 1.04 \\ R2 &= \frac{1 - \mathcal{P}^{DB}(s_{13} = .15)}{1 - \mathcal{P}^{SM}(s_{13} = .147)} = 1.00 , \end{aligned} \quad (25)$$

which demonstrates that using the S-Matrix formulation for $L=1.9\text{ km}$ and $E \simeq 4.0\text{ MeV}$ one would extract $s_{13} = .147$ from the data for which the older formalism finds $s_{13} = .15$. This is a 2% correction. We use the notation $E \simeq 4.0\text{ MeV}$ as there is uncertainty in the antineutrino energy.




Figure 3: $\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e)$. For $s_{13} = .15$ (a) $\mathcal{P}^{DB}$ and (b) $\mathcal{P}^{SM}$

Fig. 3 is the same as Fig. 2, except the baseline is $L=10$ km, as future might use a longer baseline for a larger effect given s_{13} . For $E \simeq 4.0$ MeV and $L=10$ km the ratios are

$$\begin{aligned} R1 &= \frac{1 - \mathcal{P}^{DB}(s_{13} = .15)}{1 - \mathcal{P}^{SM}(s_{13} = .15)} = 1.47 \\ R2 &= \frac{1 - \mathcal{P}^{DB}(s_{13} = .15)}{1 - \mathcal{P}^{SM}(s_{13} = .095)} = 1.00 . \end{aligned} \quad (26)$$

Thus using the S-Matrix formulation for $L=10$ km and $E \simeq 4.0$ MeV one would extract $s_{13} = .095$ from the data for which the older formalism finds $s_{13} = .15$. This is a 35% correction.

We have carried out similar calculations for the T2K project, with $E=0.6$ GeV, $L=295$ km[18]. With both a larger L and larger E than Daya Bay, we find a correction of 2.4%.

It is also important to note that our SM method gives $\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e) \neq 1.0$ even for $s_{13}=0$.

4 CP Violation $\Delta\mathcal{P}_{\mu e}^{CP}$

In this section we shall extend the derivation of the transition probability $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ of the previous section to derive the CPV probability

$$\begin{aligned} \Delta\mathcal{P}_{\mu e}^{CP} &= \mathcal{P}(\nu_\mu \rightarrow \nu_e) - \mathcal{P}(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) \\ &= |S_{12}|^2 - |\bar{S}_{12}|^2 \end{aligned} \quad (27)$$

with S_{12} defined in Eq(8) and

$$\bar{S}_{12} = c_{23}\bar{\beta} - is_{23}ae^{i\delta_{CP}}\bar{A}, \quad (28)$$

with $\bar{\beta} = \beta(V \rightarrow -V)$ and $\bar{A} = A(V \rightarrow -V)$. For example (see Eqs(16,18)) $2\bar{\omega} = \sqrt{\delta^2 + V^2 + 2\delta V \cos(2\theta_{12})}$ and $\bar{\Delta} = \Delta + (V - \delta)/2$. Using conservation of probabiltiy[31], $|A|^2 = |\bar{A}|^2$.

$$\Delta\mathcal{P}_{\mu e}^{CP} = c_{23}^2(|\beta|^2 - |\bar{\beta}|^2) - 2c_{23}s_{23}ai(Im[\beta e^{i\delta_{CP}} A^*] - Im[\bar{\beta} e^{i\delta_{CP}} \bar{A}^*]) \quad (29)$$

From Eq(29), the definitions in the previous section, defining $s \equiv \sin(\omega L)$, $c \equiv \cos(\omega L)$ one finds

$$\Delta\mathcal{P}_{\mu e}^{CP} = c_{23}^2 s_{12}^2 c_{12}^2 \delta^2 \left(\frac{s^2}{\omega^2} - \frac{\bar{s}^2}{\bar{\omega}^2} \right) + 2c_{23}s_{23}s_{12}c_{12}s_{13}\delta(\Delta - \delta s_{12}^2) \quad (30)$$

$$\begin{aligned} & \left(\sin\theta_{CP} \left(\frac{s}{\omega} (c - \cos\bar{\Delta}L) \frac{\bar{\Delta} - \omega\cos 2\theta}{\bar{\Delta}^2 - \omega^2} + \frac{\bar{s}}{\bar{\omega}} (\bar{c} - \cos\bar{\bar{\Delta}}L) \frac{\bar{\bar{\Delta}} - \bar{\omega}\cos 2\bar{\theta}}{\bar{\bar{\Delta}}^2 - \bar{\omega}^2} \right) \right. \\ & \left. - \cos\theta_{CP} \left(\frac{s}{\omega} \frac{\sin\bar{\Delta}L(\bar{\Delta} - \omega\cos 2\theta) + \sin\omega L(\omega + \bar{\Delta}\cos 2\theta)}{\bar{\Delta}^2 - \omega^2} \right. \right. \end{aligned} \quad (31)$$

$$\left. \left. - \frac{\bar{s}}{\bar{\omega}} \frac{\sin\bar{\bar{\Delta}}L(\bar{\bar{\Delta}} - \bar{\omega}\cos 2\theta) + \sin\bar{\omega}L(\bar{\omega} + \bar{\bar{\Delta}}\cos 2\theta)}{\bar{\bar{\Delta}}^2 - \bar{\omega}^2} \right) \right). \quad (32)$$

The results for $\Delta\mathcal{P}_{\mu e}^{CP}$ for $s_{13}=0.19$ are shown in Fig.4. Note that $\Delta\mathcal{P}_{\mu e}^{CP}$ depends strongly on δ_{CP} , which could lead to a measurement of this parameter. The large value of $\Delta\mathcal{P}_{\mu e}^{CP}$ for CHOOZ is promising for future experiments. $\Delta\mathcal{P}_{\mu e}^{CP}$ is so small (from about 10^{-10} to 10^{-18}) for MiniBooNE, we do not show the results. Similar results for $\Delta\mathcal{P}_{\mu e}^{CP}$ for $s_{13}=0.095$ are shown in Fig.5.




Figure 4: The ordinate is $\Delta\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ for MINOS($L=735$ km), JHF-Kamioka($L=295$ km), and CHOOZ($L=1$ km). $s_{13}=0.19$, and a, b, c, d, e for $\delta_{CP}=\pi/2, \pi/4, 0.0, -\pi/4, -\pi/2$




Figure 5: The ordinate is $\Delta\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ for MINOS($L=735$ km), JHF-Kamioka($L=295$ km), and CHOOZ($L=1$ km). $s_{13}=0.095$, and a, b, c, d, e for $\delta_{CP}=\pi/2, \pi/4, 0.0, -\pi/4, -\pi/2$

5 Conclusions

We have estimated CP violation for a variety of experimental neutrino beam facilities, for values of the parameter $s_{13}=0.19$ and $.095$, and for δ_{CP} from 90 to -90 degrees, since its value is not known. As our results show, the probability $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ is strongly dependent on s_{13} and is essentially independent of δ_{CP} (see Fig. 1), and therefore the measurement of $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$ should determine the value of the s_{13} parameter, as has been known for many years.

Our new results for $\bar{\nu}_e$ disappearance, as is being measured the Daya Bay, Double Chooz and RENO projects, however, make use of a different theoretical formulation than that used by these projects to extract s_{13} from the data. We have shown that the recent result from the Daya Bay collaboration[24] with $E \simeq 4$ MeV and $L=1.9$ km, from which it was estimated that $s_{13} \simeq .15$, by our analysis is $s_{13} \simeq .147$, a 2% correction. This is small, but the goal of these projects is 1% accuracy for s_{13} . For a baseline of $L=10$ km, with $E \simeq 4$ MeV, we find $s_{13} \simeq .097$ using the S-Matrix method, rather than $.15$, a 35% correction. Also, our SM method gives $\mathcal{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e) \neq 1.0$ even for $s_{13}=0$.

The CP violation probability (CPV), $\Delta\mathcal{P}_{\mu e}^{CP}$, is strongly dependent on both of these important parameters. After the Double Chooz/Daya Bay/RENO determination of s_{13} , both the JHF-Kamioka and Double Chooz projects might be able to determine the value of δ_{CP} , since for most of the values of δ_{CP} these projects would have nearly a 1% CPV, as shown in Figs. 4 and 5. No experiments are possible now to test CPV via neutrino oscillations, since beams of both neutrino and antineutrino with the same flavor would be needed. However, in the future such beams might be available. Our results should help in planning such future experiments.

Acknowledgements

This work was supported in part by the NSF grant PHY-00070888, in part by the DOE contracts W-7405-ENG-36 and DE-FG02-97ER41014, and in part by a grant from the Pittsburgh Foundation. We thank Prof Ma Wei-xing, IHEP Beijing, for information about the Daya Bay project.

References

- [1] J.H. Christenson, J.W. Cronin, V.L. Fitch, and R. Turlay, Phys. Rev. Lett. **13**, 138 (1964)
- [2] J.W. Cronin, P.F. Kunz, W.S. Risk, and P.C. Wheeler, Phys. Rev. Lett. **18**, 25 (1967)
- [3] L.S. Littenberg, Phys. Rev. D **39**, 3322 (1989)

- [4] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973); L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983)
- [5] A.J. Buras and S. Uhlig, Rev. Mod. Phys. **90**, 965 (2008)
- [6] M. Benzke, S.J. Lee, M. Neubert, and G. Paz, Phys. Rev. Lett. **106**, 141801 (2011)
- [7] The MiniBooNE Collaboration, Phys. Rev. Lett. **105**, 181801 (2010)
- [8] The JHF-Kamioka neutrino Project, arXiv:hep-ex/106019
- [9] The Minos Collaboration, Phys. Rev. Lett. **103**, 261802 (2009); Phys. Rev. **D 81**, 052004 (2010)
- [10] CHOOZ Collaboration, M. Apollonio *et al* Eur. J. **C 27**, 331 (2003)
- [11] Double Chooz Collaboration, F. Ardellier *et. al.* hep-ex/0606025 (2006)
- [12] H. Davoudiasl, H-S. Lee, and W. J. Marciano, Phys. Rev. **D 84**, 013009 (2011)
- [13] M.C. Gonzalez-Garcia, M. Maltoni, and J. Salvado, arXiv:1103.4365/hep-ph; JHEP 1105: 075 (2011)
- [14] The ISS Working Group, arXiv:0710.4947/hep-ph
- [15] LBNE Project, lbne.fnal.gov/project/general-info.shtml
- [16] V. Barger *et al*, Report of the US long baseline neutrino experiment study, arXiv:0705.4396/hep-ex
- [17] L.S. Kisslinger, arXiv:1108.4062/hep-ph
- [18] K. Abe *et al*, T2K Collaboration, Phys. Rev. Lett **107**, 041801 (2011)
- [19] Daya Bay Collaboration, arXiv:0701029/hep-ex (2007)
- [20] Daya Bay Collaboration, C-J Lin, arXiv:1101.0261/hep-ex (2011)
- [21] M. Kuze for Double Chooz Collaboration, arXiv:1109.0074/hep-ex (2011)
- [22] P. Novella, arXiv:1105.6079/hep-ex
- [23] RENO Collaboration, J.K. Ahn, *et al*, arXiv:1003.1391/hep-ex (2010)
- [24] Daya Bay Collaboration, [http : //dayawane.ihep.ac.cn/docs/DYB_{rate_prl_APS}.pdf](http://dayawane.ihep.ac.cn/docs/DYB_rate_prl_APS.pdf)

- [25] J. Arafune and J. Sato, Phys. Rev. **D 55**, 1653 (1997) fnal
- [26] S.M Bilenky, C. Giunti, and W. Grimus, Phys. Rev.**D 58**, 033001 (1997)
- [27] K. Kimura, A. Takamura, and H. Yokomakura, Phys. Rev. **D 66**, 073005 (2002)
- [28] E.M. Henley, M.B. Johnson, and L.S. Kisslinger, arXiv:hep-ph/1102.5106; Int J. Mod. Phys. E **20**, 2463 (2012)
- [29] E. Akhmedov, P. Huber, M. Lindner, and T. Ohlsson, Nucl. Phys. **B608**, 394 (2001)
- [30] M. Freund, Phys. Rev. **D64**, 053003 (2001)
- [31] M. Jacobson and T. Ohlsson, Phys. Rev. **D 69**, 013003 (2004)